

Thermal self-energies using light-front quantization

H. Arthur Weldon

Department of Physics, West Virginia University, Morgantown, West Virginia 26506-6315

(Dated: February 7, 2008)

A recent paper by Alves, Das, and Perez contains a calculation of the one-loop self-energy in ϕ^3 field theory at $T \neq 0$ using light-front quantization and concludes that the self-energy is different than the conventional answer and is not rotationally invariant. The changes of variable displayed below show that despite the complicated appearance of the thermal self-energy in light-front variables, it is exactly the same as the conventional result.

In Ref. [1], Alves, Das, and Perez introduce the technique of light-front quantization into thermal field theory using a heat bath at rest. As shown in [2] the appropriate quantization evolves the system in the $x^+ = (x^0 + x^3)/\sqrt{2}$ coordinate while keeping constant x^1, x^2 , and x^3 . (Normal light-front quantization keeps x^1, x^2 , and $x^- = (x^0 - x^3)/\sqrt{2}$ constant.) The momenta conjugate to x^+, x^1, x^2, x^3 are k^0, k^1, k^2, k^+ as can be seen from the identity

$$k^0 x^0 - k^3 x^3 - \mathbf{k}_\perp \cdot \mathbf{x}_\perp = \sqrt{2} k^0 x^+ - \sqrt{2} k^+ x^3 - \mathbf{k}_\perp \cdot \mathbf{x}_\perp.$$

In the imaginary time propagator, x^+ is made negative imaginary: $-i\beta \leq \sqrt{2}x^+ \leq 0$. In the Fourier transform of the propagator $k^0 = i2\pi nT$ whereas k^+ and $\mathbf{k}_\perp$ are real. The relation

$$(k^0)^2 - (k^3)^2 - k_\perp^2 = 2\sqrt{2}k^0 k^+ - 2(k^+)^2 - k_\perp^2$$

immediately leads to the propagator

$$G(k^+, k_\perp, n) = \frac{1}{i4\sqrt{2}\pi n k^+ - 2(k^+)^2 - \omega_k^2},$$

where $\omega_k^2 = k_\perp^2 + m^2$ is the transverse energy.

One of the interesting calculations performed by Alves, Das, and Perez [1] using this propagator is the one-loop self-energy for a scalar field theory with self-interaction $g\phi^3/3!$. The result of performing the summation over the loop integer n is given in Eq. (40) of Ref. [1]:

$$\Pi(p) = \frac{g^2}{8} \int \frac{dk^+ d^2 k_\perp}{(2\pi)^3} \frac{\coth(X_1/2T) - \coth(X_2/2T)}{Y}. \quad (1)$$

During the summation the external variable p^0 is an integer multiple of $2\pi iT$, but after the summation p^0 is continued to real values. The quantities X_1, X_2 , and Y are complicated functions of the integration variables $k^+, \mathbf{k}_\perp$ and of the external variables $p^0, p^+, \mathbf{p}_\perp$:

$$\begin{aligned} X_1 &= \frac{\omega_k^2 + 2(k^+)^2}{2\sqrt{2}k^+} \\ X_2 &= \frac{\omega_{k+p}^2 + 2(k^+ + p^+)^2}{2\sqrt{2}(k^+ + p^+)} \\ Y &= 2\sqrt{2}k^+(k^+ + p^+)[-X_1 + X_2 - p^0]. \end{aligned}$$

The self-energy (1) looks quite different from the usual result and is not manifestly invariant under $O(3)$ rotations of the external momentum $\mathbf{p}$.

The following will describe a change of integration variable from k^+ to a new variable k^3 that is chosen to make the self-energy a function of the two variables p^0 and $\mathbf{p}^2 = p_\perp^2 + (\sqrt{2}p^+ - p^0)^2$. The final answer is the sum of Eqs. (2), (3), (4), and (5).

(1a) For the term $\cosh(X_1/2T)/Y$ in Eq. (1), when k^+ is positive change to a new integration variable k^3 defined by

$$k^+ = \frac{1}{\sqrt{2}} \left[k^3 + \sqrt{m^2 + k_\perp^2 + (k^3)^2} \right].$$

The range of k^3 is $-\infty \leq k^3 \leq \infty$. The Jacobian of the transformation is $dk^+/dk^3 = k^+/E_k$, where $E_k = \sqrt{m^2 + k^2}$ is the square root displayed above. Under this change,

$$X_1 = E_k; \quad Y = k^+ [(E_{k+p})^2 - (p^0 + E_k)^2],$$

where $E_{k+p} = \sqrt{m^2 + (\mathbf{k} + \mathbf{p})^2}$. The factor k^+ in the Jacobian cancels a similar factor in Y and yields a contribution to the self-energy

$$\Pi_{1a} = \frac{g^2}{8} \int \frac{d^2 k_\perp}{(2\pi)^3} \int_{-\infty}^{\infty} \frac{dk^3}{E_k} \frac{\coth(E_k/2T)}{(E_{k+p})^2 - (p^0 + E_k)^2}. \quad (2)$$

This integrand is invariant under simultaneous rotations of the vectors $\mathbf{k}$ and $\mathbf{p}$. Thus Π_{1a} depends only on $|\mathbf{p}|$ and p^0 .

(1b) For term $\cosh(X_2/2T)/Y$ in Eq. (1), when k^+ is negative make the change of variable

$$k^+ = \frac{1}{\sqrt{2}} \left[k^3 - \sqrt{m^2 + k_\perp^2 + (k^3)^2} \right],$$

where $-\infty \leq k^3 \leq \infty$. The Jacobian of the transformation is $dk^+/dk^3 = -k^+/E_k$ and

$$X_1 = -E_k; \quad Y = k^+ [(E_{k+p})^2 - (p^0 - E_k)^2].$$

The corresponding self-energy contribution is

$$\Pi_{1b} = \frac{g^2}{8} \int \frac{d^2 k_\perp}{(2\pi)^3} \int_{-\infty}^{\infty} \frac{dk^3}{E_k} \frac{\coth(E_k/2T)}{(E_{k+p})^2 - (p^0 - E_k)^2}. \quad (3)$$

The sum of Eqs. (2) and (3) is an even function of p^0 .

(2a) In the second term in Eq. (1), $\cosh(X_2/2T)/Y$, when $k^+ + p^+ > 0$ then change to k^3 given by

$$k^+ = \frac{1}{\sqrt{2}} \left[k^3 - p^0 + \sqrt{m^2 + (\mathbf{k}_\perp + \mathbf{p}_\perp)^2 + (k^3 + p^3)^2} \right],$$

where $-\infty \leq k^3 \leq \infty$. Using $dk^+/dk^3 = (k^+ + p^+)/E_{k+p}$, and

$$X_2 = E_{k+p}; \quad Y = (k^+ + p^+)[(p^0 - E_{k+p})^2 - E_k^2].$$

This contribution is

$$\Pi_{2a} = \frac{g^2}{8} \int \frac{d^2 k_\perp}{(2\pi)^3} \int_{-\infty}^{\infty} \frac{dk^3}{E_{k+p}} \frac{\coth(E_{k+p}/2T)}{E_k^2 - (p^0 - E_{k+p})^2}. \quad (4)$$

(2b) In the second term in Eq. (1), $\cosh(X_2/2T)/Y$, if $k^+ + p^+ < 0$ make the change of variable

$$k^+ = \frac{1}{\sqrt{2}} \left[k^3 - p^0 - \sqrt{m^2 + (\mathbf{k}_\perp + \mathbf{p}_\perp)^2 + (k^3 + p^3)^2} \right],$$

where $-\infty \leq k^3 \leq \infty$. Since $dk^+/dk^3 = -(k^+ + p^+)/E_{k+p}$, and

$$X_2 = -E_{k+p}, \quad Y = (k^+ + p^+)[(p^0 + E_{k+p})^2 - E_k^2],$$

the contribution to the self-energy is

$$\Pi_{2b} = \frac{g^2}{8} \int \frac{d^2 k_\perp}{(2\pi)^3} \int_{-\infty}^{\infty} \frac{dk^3}{E_{k+p}} \frac{\coth(E_{k+p}/2T)}{E_k^2 - (p^0 + E_{k+p})^2}. \quad (5)$$

The sum of Eq. (4) and (5) is an even function of p^0 .

The sum of these four contributions Eq. (2), (3), (4), and (5) is the standard answer for the self-energy [3]. Therefore the light-front formulation is a different, and in some cases a more efficient [2], way of organizing the calculation, but the results are the same.

Acknowledgments

This work was supported in part by the U.S. National Science Foundation under Grant No. PHY-0099380.

-
- [1] V.S. Alves, Ashok Das, and Silvana Perez, Phys. Rev. D **66**, 125008 (2002).
 [2] H.A. Weldon, Phys. Rev. D **67**, 0850XX (2003).

- [3] A. Das, *Finite Temperature Field Theory* (Cambridge, University Press, Cambridge, England, 1996), page 24.